

# Is the Higgs a Composite Scalar?

Evan Weinberg

Boston University, Boston, MA

*weinbe2@bu.edu*

March 19, 2016



RBRC Workshop on Lattice Gauge Theories 2016

9-11 March 2016 *Brookhaven National Laboratory, Physics Department*

Based on work in [arXiv:1601.04027](https://arxiv.org/abs/1601.04027)



## Lattice **S**trong **D**ynamics Collaboration



Xiao-Yong Jin  
James Osborn



Joe Kiskis



Rich Brower  
Claudio Rebbi  
Evan Weinberg



Oliver Witzel



Anna Hasenfratz  
Ethan Neil



Enrico Rinaldi  
Pavlos Vranas



Ethan Neil



David Schaich



Meifeng Lin



Tom Appelquist  
George Fleming  
Andy Gasbarro



Graham Kribs

1

# The Higgs Boson




Figure: [Phys. Lett. B 716 (2012)]




Figure: [Eur.Phys.J. C75 (2015)]

- The Higgs Boson looks very Standard Model.
- There's still the need for a UV completion.

# A Composite UV Completion





Figure: Modified from Wikipedia: "Standard Model"

- Higgs:  $\bar{Q} Q$  scalar composite of strong dynamics.

# Possibly Composite Teases from the LHC



- 750 GeV diphoton bump:  $\eta'$ -like state from a new composite sector?
- 2 TeV diboson bump:  $\rho$ -like state, not fully excluded, especially if state is *broad*.

Figure: ATLAS Diphoton @ 13 TeV

[ATLAS-CONF-2015-081]

The new sector can't just be scaled up QCD.

- QCD has...
  - a broad scalar close to the vector mass,
  - a large S-parameter, and
  - no walking regime.
- More flavors can produce different, interesting behavior.




Figure: [Phys.Rev. D76 (2007)]

- It is not easy to build a viable composite model for EWSB.
- Broadly speaking, there are two steps:
  - 1 Pick a general model and study if it has certain features.
  - 2 Worry about connecting it to the Standard Model.

- It is not easy to build a viable composite model for EWSB.
- Broadly speaking, there are two steps:
  - 1 Pick a general model and study if it has certain features. ← This talk
  - 2 Worry about connecting it to the Standard Model.



# Multi-flavor QCD





- The interest in multi-flavor QCD is motivated by the beta function.

$$\beta(g) = -\beta_0 g^3 - \beta_1 g^5 + \mathcal{O}(g^7)$$

$$\beta_0 = \left[ \frac{11}{3} N_c - \frac{2}{3} N_f \right] / (4\pi)^2$$


$$\beta_1 = \left[ \frac{34}{3} N_c^2 - \left( \frac{13}{3} N_c - \frac{1}{N_c} \right) N_f \right] / (4\pi)^4$$

$$\beta_1 = 0 \rightarrow N_f \approx 8.05$$




# The lattice and many-fermion physics

- We can use the lattice as a probe of **new** non-perturbative physics.
- The lattice has indicated S parameter suppression with more flavors.
- It has also indicated that  $SU(3)$  12 flavor is conformal.



[Schaich, LATTICE 2011, arXiv:1111.4993]



[Aoki et al., LATTICE 2014, arXiv:1501.06660]

- And, in the mass-deformed theory, has a light scalar.

- ❶ Pick a general model and study if it has certain features.
  - Lattice study of  $SU(3)$  with 8 fundamental flavors
    - Chiral Kogut-Susskind “Staggered” fermions: multiples of 4 flavors.
  - Right near 2-loop (strongly coupled...) opening of conformal window.
  - Likely confining, but possibly conformal. We cannot definitively tell.
    - “Feature, not a bug.”
- ❷ Worry about connecting it to the Standard Model.

## Aside: Partner 4+8 Project

- We have a partner project with 4 chiral, 8 massive flavors.
  - Brower et al., “A composite Higgs model at a conformal fixed point”
  - Available at [arXiv:1512.02576](https://arxiv.org/abs/1512.02576).
- Begin to address the question: “What happens when we decouple some fermions in a (near-)conformal theory?”
  - Important for non-minimal ( $>3$  pNGB) composite models.




- $SU(3)$  gauge group, 8 continuum fermions
- Gauge action: fundamental-adjoint with  $\beta_a = -\beta/4$  [Cheng et al. 2013][Cheng et al. 2014]
  - Negative adjoint term helps avoid lattice fixed point
- Fermion action: 1-nHYP smeared staggered [Hasenfratz et al. 2007]
  - nHYP smearing helps push to stronger coupling (avoid lattice phase)
- Software: HMC and most measurements in FUEL [J. Osborn]
- Runs:
  - “Fixed”  $M_\pi L$ :
  - $24^3, m = 0.00889$ ;  $32^3, m = 0.005$ ;  $48^3, m = 0.00222$ ;  $64^3, m = 0.00125$
  - Finite volume:
  - $24^3$  and  $32^3, m = 0.0075$
  - Large mass (connect to LatKMI):
  - $24^3, m = 0.035$

# Quark-Line Connected Spectrum


- Interested in ground state in many isomultiplet channels
- Using QCD language:
  - $\pi (0^{-+})$  —  $W^{\pm}, Z$
  - $\rho (1^{--})$  — di-boson resonance?
  - $N (\frac{1}{2}^{+})$  — dark matter?
  - Use wall sources to project onto ground states [Gupta et al. 1991]
  - Also interested in decay constants (93 MeV convention)
    - $F_{\pi}$  — electroweak symmetry vev  $\approx 250$  GeV
    - $F_{\rho}, F_{a_1}$  — production cross-sections, chiral restoration?
    - Use current-current correlators to avoid renormalization
  - $a_0 (0^{++})$  — detector signal?
  - $a_1 (1^{++})$  — detector signal?

# Non-singlet spectrum



- Unlike QCD: shouldn't trust ChiPT here.
- Spectrum depends heavily on the fermion mass.
  - Curvature at small  $m_q$

# Chirally broken?



- Looks chirally broken, or conformal with scaling corrections?
  - Boulder 12 flavor results on conformal scaling corrections:  
[Cheng et al. 2014]
- Interesting dynamics:  $M_\rho > 2M_\pi$



# $0^{++}$ scalar correlation function

$$\text{Diagram 1} - \text{Diagram 2} \Rightarrow \int_0^2 e^{-m_\sigma^2 t} + \dots$$

Isosinglet Scalar ( $0^{++}$ )

$$S(t) = \langle \bar{\psi}_a \psi_a(0) \bar{\psi}_a \psi_a(t) \rangle$$

$$\begin{aligned} &= \frac{1}{Z} \int [dU d\bar{\psi} d\psi] \left\{ 2 \overline{\bar{\psi}(0)\psi(0)} \overline{\bar{\psi}(t)\psi(t)} - \overline{\bar{\psi}(0)\psi(0)} \overline{\bar{\psi}(t)\psi(t)} \right\} e^{-\frac{1}{g^2} F^2 - \bar{\psi}_i \not{D} \psi_i - m_q \bar{\psi}_i \psi_i} \\ &= \frac{1}{Z} \int [dU] \left\{ 2 G_F(0,0) G_F(t,t) - (G_F(0,t) \gamma_5)^2 \right\} \det(D^\dagger D + m_q^2)^{N/2} e^{-\frac{1}{g^2} F^2} \\ &\equiv 2D(t) - C(t) \end{aligned}$$

- Factor  $G_F(t, t)$  means we need every point to itself.

# Strategy for disconnected diagrams

- Fitting the isosinglet,  $0^{++}$  meson requires disconnected diagrams.
- 6  $U(1)$  sources with dilution in time, color, and even/odd spatially
- Improved estimator for  $\langle \bar{\psi}\psi \rangle$
- Still need large statistics to suppress gauge noise

What this gets us:

$$C(t) = -A_{a_0} e^{-M_{a_0} t} - (-1)^t \left( A_{\pi_{sc}} e^{-M_{\pi_{sc}} t} \right) + \dots$$

$$S(t) \equiv 2D(t) - C(t) = A_{\sigma} e^{-M_{\sigma} t} + (-1)^t \left( A_{\eta} e^{-M_{\eta} t} \right) + \dots$$

↓ [Aoki et al., LATTICE 2014, arXiv:1501.06660]

$$2D(t) = A_{\sigma} e^{-M_{\sigma} t} - A_{a_0} e^{-M_{a_0} t} + (-1)^t \left( A_{\eta} e^{-M_{\eta} t} - A_{\pi_{sc}} e^{-M_{\pi_{sc}} t} \right) + \dots$$


# Strategy for disconnected diagrams, continued

$$S(t) \equiv 2D(t) - C(t) = A_\sigma e^{-M_\sigma t} + (-1)^t \left( A_\eta e^{-M_\eta t} \right) + \dots$$



$$2D(t) = A_\sigma e^{-M_\sigma t} - A_{a_0} e^{-M_{a_0} t} + (-1)^t \left( A_\eta e^{-M_\eta t} - A_{\pi_{sc}} e^{-M_{\pi_{sc}} t} \right) + \dots$$

If  $M_{0^{++}} < M_{a_0}$ , we can extract from both  $D(t)$ ,  $S(t)$  at large  $t$ .

- Correlated fit to both parity states
- **Vacuum subtraction** introduces large uncertainties
- Fit the finite difference:
  - $D(t+1) - D(t)$
  - $S(t+1) - S(t)$
- Average resulting  $M_{0^{++}}$ , errors in quadrature




# Singlet spectrum





- Generally, the  $0^{++}$  tracks the Goldstone boson in this regime.
  - LatKMI results: [Y. Aoki et al. 2013] [Y. Aoki et al. SCGT15] (Thank you!)
- Very different from QCD!

# Spectrum Overview




- Look at ratios with  $F_\pi \approx v \approx 250 \text{ GeV}$
- Light  $0^{++} \approx \pi < \rho, 2\pi$
- $M_{0^{++}}/F_\pi \approx 4$ : What would a top loop do?
- Rich spectrum of other states.
- $M_\rho/F_\pi \approx 8$ : 2TeV di-boson resonance?
  - Ratio seen in QCD
  - Also seen in  $SU(3)$  2 flavor sextet [LatHC 2015 (LATTICE2015)]
  - Perhaps a general feature?

# Other Results: Decay Constants



- Decay constants extracted from current correlators: no “Z” factor.
- $F_\pi$  is smallest;  $F_\rho$ ,  $F_{a_1}$  degenerate w/in errors.
- Interesting: Check Kawarabayashi-Suzuki-Riazuddin-Fayyazuddin (KSRF) relations:  $F_\rho M_\rho = 2F_\pi^2 g_{\rho\pi\pi}$ ;  $M_\rho^2 = 2F_\pi^2 g_{\rho\pi\pi}^2$ .
  - Predicts  $F_\rho = \sqrt{2}F_\pi$ —we (roughly) see that!

# Vector Meson Phenomenology




- Production rate related to  $F_\rho$ .
- Decay via longitudinal coupling to Goldstones ( $W^\pm, Z$ ):  $g_{\rho\pi\pi}$
- Estimate  $g_{\rho\pi\pi}$  via KSRF:  $g_{\rho\pi\pi} = \frac{M_\rho}{\sqrt{2}F_\pi}$ .
- Estimate  $\Gamma_\rho \approx \frac{g_{\rho\pi\pi}^2 M_\rho}{48\pi} \approx \frac{M_\rho^3}{96\pi F_\pi^2}$  (Assuming  $M_\rho \gg M_\pi$ )
  - Using  $M_\rho \approx 2\text{ TeV}$ ,  $F_\pi \approx 250\text{ GeV} \rightarrow \Gamma_\rho \approx 450\text{ GeV}$ .

- ① Pick a general model and study if it has certain features.
  - 8 flavors is a great model to learn about light scalar dynamics.
  - What is the low energy theory when there's a light scalar?
    - We're on the UV complete lattice.
    - We can compute  $\pi$ - $\pi$  scattering,  $\pi$ - $\sigma$  scattering,  $\sigma$ - $\sigma$  scattering.
    - From a field theory standpoint, we can learn a lot.
- ② Worry about connecting it to the Standard Model.



# Upcoming work


$I = 2 \pi^+ \pi^+$  scattering



$\rightarrow W^+ W^+$  scattering

---

Scalar form factor of the Pion




$\rightarrow$  Higgs absorption/emission by  $W^\pm, Z$

- ① Pick a general model and study if it has certain features.
  - 8 flavors is a great model to learn about light scalar dynamics.
  - What is the **low energy theory** when there's a light scalar?
    - We're on the UV complete lattice.
    - We can compute  $\pi$ - $\pi$  scattering,  $\pi$ - $\sigma$  scattering,  $\sigma$ - $\sigma$  scattering.
    - From a field theory standpoint, we can learn a lot.
- ② Worry about connecting it to the Standard Model.

Thank you!

# Backup


# Backup: The Lattice



- Studies are done on 4-dimensional  $L \times L \times L \times (2L)$  lattices.
- Common values are  $L = 24, 32 \rightarrow \mathcal{O}(1 \text{ million})$  sites.

$$\mathcal{Z} = \int [dU d\bar{\psi} d\psi] e^{-\frac{1}{g^2} F^2 - \bar{\psi}_i \not{D} \psi_i - m_\ell \bar{\psi}_\ell \psi_\ell - m_h \bar{\psi}_h \psi_h}$$

# Backup: The Lattice



- Studies are done on 4-dimensional  $L \times L \times L \times (2L)$  lattices.
- Common values are  $L = 24, 32 \rightarrow \mathcal{O}(1 \text{ million})$  sites.

$$\mathcal{Z} = \int [dU] \det(D^\dagger D + m_h^2)^{N_h/2} \det(D^\dagger D + m_\ell^2)^{N_\ell/2} e^{-\frac{1}{g^2} F^2}$$

- Having multiple flavors is just adding more fermion determinants.

# Backup: 8 flavors finite temperature studies

- We base our 8 flavor runs on existing results.




Figure: Finite T studies by Boulder / LSD, in preparation

- Run at strong couplings safe from deconfinement and lattice phases.

- Lattice study of  $SU(3)$  with 8 fundamental flavors
  - Gauge action: fundamental-adjoint with  $\beta_a = -\beta/4$  [Cheng et al. 2013][Cheng et al. 2014]
  - Fermion action: nHYP smeared staggered [Hasenfratz et al. 2007]
  - Software: HMC and most measurements in FUEL [J. Osborn]



# Backup: Setting and varying a scale

- Along the gradient flow...  
[arXiv:1006.4518]

$t$  = gradient flow time

- ...look at a quantity...

$$g_{WF}^2 \left( \mu = \frac{1}{\sqrt{8t}} \right) = \frac{1}{\mathcal{N}} \langle t^2 E(t) \rangle$$

- ...relative to a consistent IR scale.

$$g_{WF}^2(\mu_0 = \frac{1}{\sqrt{8t_0}}) = \frac{0.3}{\mathcal{N}}$$

- $\mu_0$  is an energy scale.
- $\sqrt{8t_0}$  is a length scale.




Figure:  $48^3 \times 96$ ,  $m_q = 0.00222$

- There are five (maybe 6) isosinglet scalars below the charm threshold.
- $f_0(500)$ ,  $f_0(980)$ ,  $f_0(1370)$ ,  $f_0(1500)$ ,  $f_0(1710)$  (and maybe  $f_0(1790)$ ).
- Quark model: only two can be predominantly  $\bar{Q}Q$ .
- Others: meson molecule, diquark pair, glueballs?

## Backup: KSRF Relations

$$F_\rho M_\rho = 2F_\pi^2 g_{\rho\pi\pi}$$

$$M_\rho^2 = 2F_\pi^2 g_{\rho\pi\pi}^2$$

These imply:

$$F_V = \sqrt{2}F_\pi$$

$$g_{\rho\pi\pi} = M_\rho/F_\rho$$

These are good at  $\approx 20\%$  in QCD, even at heavier pion masses.

Need to add links to citations: use in QCD, how it comes from simple models of massive vectors.

# Backup: Strategy for disconnected diagrams

- 8 flavors uses the following setup!
  - 6  $U(1)$  sources with dilution in time, color, and even/odd spatially
  - Improved estimator for  $\langle \bar{\psi}\psi \rangle$
  - Dilution in time, color, even/odd space
  - Improved estimator for disconnected piece
  - Still need large statistics to suppress gauge noise
- Analysis strategy.
  - Correlated fit to both parity states
  - **Vacuum subtraction** introduces large uncertainties
  - Fit an additional constant
  - Equivalent to fitting the finite difference  $C(t+1) - C(t)$

$$C(t) = c_{0++} \cosh(M_{0++}(T/2 - t)) + c_{\tilde{\pi}_{sc}} (-1)^t \cosh(M_{\tilde{\pi}_{sc}}(T/2 - t)) + v$$